

THE MODULUS OF WHITTAKER FUNCTIONS

HANS VOLKMER

ABSTRACT. The paper discusses some properties of the modulus $|W_{k,m}(z)|$ of the Whittaker function $W_{k,m}(z)$. In particular, completely monotone functions expressed in terms of $|W_{k,m}(z)|$ are found. The results follow from an integral representation for products of Whittaker functions due to Erdélyi (1938).

1. INTRODUCTION

Recently, Boris Belinskiy asked the author the following question: Consider the quotient of Whittaker functions

$$(1.1) \quad \frac{W_{k,m}(\tau z)}{W_{k,m}(z)},$$

where k is purely imaginary, m is real, $|\arg z| < \frac{1}{2}\pi$, $\tau > 1$. Show that the quotient (1.1) is bounded for fixed z and τ as a function of k, m . This problem occurs in Belinskiy's work in scattering theory. A straightforward approach to this problem is to investigate the asymptotic behavior of the Whittaker function $W_{ir,m}$ as the real variables r, m tend to infinity. However, this approach faces some difficulties. It appears that the required asymptotic results are not available. The handbook of mathematical functions [3, Chapter 13] does give asymptotic results where both k and m tend to infinity but they do not include imaginary k . Olver [4, page 401] has such results but only for fixed m . Moreover, it will be difficult to obtain explicit bounds for (1.1) by using asymptotic methods.

We show in this paper that the quotient (1.1) can be estimated in a very simple way using an integral representation for products of Whittaker functions due to Erdélyi [2]. We also show that the function

$$t \mapsto t^{-1} e^{t\Re x} |W_{k,m}(tx)|^2, \quad t > 0$$

is completely monotone for appropriate values of k, m and x .

Erdélyi's formula is known for a long time. However, the conclusions we draw from it appear to be new.

2010 *Mathematics Subject Classification.* 33C15, 33C10.

Key words and phrases. Whittaker functions; Bessel functions; completely monotone functions.

2. ERDÉLYI'S INTEGRAL FORMULA

Erdélyi [2] used the convolution theorem for the Laplace transform to derive the following integral representation for products of Whittaker functions.

Theorem 2.1. *Suppose $x, y \in \mathbb{C}$, $|\arg x| < \pi$, $|\arg y| < \pi$, $t > 0$, $m \in \mathbb{C}$, $k, l \in \mathbb{C}$ with $\Re(1 - k - l) > 0$. Then we have*

$$(2.1) \quad \begin{aligned} & (t^2 xy)^{-1/2} e^{\frac{1}{2}t(x+y)} W_{k,m}(tx) W_{l,m}(ty) \\ &= \frac{(xy)^m}{\Gamma(1 - k - l)} \int_0^\infty e^{-tu} (x+u)^{k-\frac{1}{2}-m} (y+u)^{l-\frac{1}{2}-m} u^{-k-l} \\ & \quad \times {}_2F_1\left(\frac{1}{2} + m - k, \frac{1}{2} + m - l; 1 - k - l; \frac{u(x+y+u)}{(x+u)(y+u)}\right) du. \end{aligned}$$

Erdélyi [2, (1) on page 873] has (2.1) with $t = 1$. Buchholz [1, (10) on page 89] has (2.1) including t . It is easy to see that we can obtain the formula with t from the special case $t = 1$. For our purpose, we need (2.1) including t . The function ${}_2F_1$ is the hypergeometric function. If $u = 0$ we choose the principal value of ${}_2F_1$ and analytically continue this branch along the path of integration. The powers appearing in (2.1) are assigned their principal values.

If m is real, $y = \bar{x}$, $l = \bar{k}$, we obtain the following result as a special case.

Corollary 2.2. *Suppose $x \in \mathbb{C}$, $|\arg x| < \pi$, $t > 0$, $m \in \mathbb{R}$, $k \in \mathbb{C}$, $\Re k < \frac{1}{2}$. Then we have*

$$(2.2) \quad \begin{aligned} & (t|x|)^{-1} e^{t\Re x} |W_{k,m}(tx)|^2 \\ &= \frac{|x|^{2m}}{\Gamma(1 - 2\Re k)} \int_0^\infty e^{-tu} \left| (x+u)^{k-\frac{1}{2}-m} \right|^2 u^{-2\Re k} \\ & \quad \times {}_2F_1\left(\frac{1}{2} + m - k, \frac{1}{2} + m - \bar{k}; 1 - 2\Re k; \frac{u(2\Re x + u)}{|x+u|^2}\right) du. \end{aligned}$$

The argument $\frac{u(2\Re x + u)}{|x+u|^2}$ of the hypergeometric function in (2.2) is real and less than 1.

3. CONSEQUENCES

We use Corollary 2.2 to estimate (1.1).

Theorem 3.1. *Suppose $x \in \mathbb{C}$, $|\arg x| \leq \frac{1}{2}\pi$, $m \in \mathbb{R}$, $k \in \mathbb{C}$, $\Re k < \frac{1}{2}$ and $\tau \geq 1$. Then*

$$(3.1) \quad \left| \frac{W_{k,m}(\tau x)}{W_{k,m}(x)} \right|^2 \leq \tau e^{(1-\tau)\Re x}.$$

Proof. Since $\Re x \geq 0$, $\frac{u(2\Re x + u)}{|x+u|^2}$ lies between 0 and 1. Also, the hypergeometric power series appearing in (2.2) has nonnegative coefficients. Therefore,

the right-hand side of (2.2) is a decreasing function of t , and we obtain, for $t = \tau \geq 1$,

$$(\tau|x|)^{-1}e^{\tau\Re x}|W_{k,m}(\tau x)|^2 \leq |x|^{-1}e^{\Re x}|W_{k,m}(x)|^2.$$

This implies (3.1) $\square$

It is important that the bound on the right-hand side of (3.1) is independent of k and m in order to answer the question raised in the introduction.

Consider the hypergeometric function

$$F(z) = {}_2F_1\left(\frac{1}{2} + m - k, \frac{1}{2} + m - \bar{k}; 1 - 2\Re k; z\right)$$

under the assumptions on k, m from Corollary 2.2. Then $F(z) > 0$ for $z \in [0, 1)$. If $F(z)$ has a zero $z < 0$ then let $p = p(k, m)$ denote the largest negative zero of $F(z)$. If there is no negative zero, set $p = -\infty$.

Theorem 3.2. *Suppose that $k \in \mathbb{C}$, $\Re k < \frac{1}{2}$, $m \in \mathbb{R}$, and $x \in \mathbb{C}$ such that $|\arg x| < \pi$ if $p = -\infty$, and otherwise $|\arg x| \leq \theta$, where θ is the angle between $\frac{1}{2}\pi$ and π for which*

$$\tan \theta = -\frac{1}{(-p(k, m))^{1/2}}.$$

Then the function

$$(3.2) \quad f(t) = t^{-1}e^{t\Re x}|W_{k,m}(tx)|^2$$

is (strictly) completely monotone on $(0, \infty)$, that is $(-1)^n f^{(n)}(t) > 0$ for all $t > 0$ and $n = 0, 1, 2, \dots$

Proof. If $|\arg x| \leq \frac{1}{2}\pi$ then (2.2) shows that $f(t)$ is the Laplace transform of a positive function, so the statement follows. If $\frac{1}{2}\pi < |\arg x| < \pi$, then the function

$$u \mapsto \frac{u(2\Re x + u)}{|x + u|^2}, \quad u \geq 0,$$

attains an absolute minimum at $u = -\Re x$ with value $-\left(\frac{\Re x}{\Im x}\right)^2$. Under the assumption $|\arg x| \leq \theta$, this minimum value is greater than or equal to $p(k, m)$. Therefore, $f(t)$ is still the Laplace transform of a positive function (which has one zero when $|\arg x| = \theta$.) $\square$

We should mention that the condition $|\arg x| \leq \theta$ is sharp. If this condition is not satisfied (with $|\arg x| < \pi$) then $f(t)$ is not completely monotone because of Bernstein's theorem [5, Thm 12b, page 161]. As a consequence of Theorem 3.2 we obtain that the function $W_{k,m}(x)$ has no zeros in the sector $|\arg x| \leq \theta$. It is clear that Theorem 3.1 could also be extended to the sector $|\arg x| \leq \theta$.

We identify a case when $p(k, m) = -\infty$.

Theorem 3.3. *Suppose that $k < \frac{1}{2}$,*

$$k - \frac{1}{2} \leq m \leq \frac{1}{2} - k$$

and $x \in \mathbb{C}$, $|\arg x| < \pi$. Then the function (3.2) is (strictly) completely monotone on $(0, \infty)$.

Proof. The integral representation [3, 15.6.1] shows that ${}_2F_1(a, b; c; z) > 0$ when $a, b, c \in \mathbb{R}$, $c > b > 0$ and $z < 1$. This remains true if either $b = 0$ or $b = c$. It follows that Theorem 3.2 applies with $p(k, m) = -\infty$. $\square$

It is of interest to look at these results for the special case of the modified Bessel function K_ν connected to Whittaker functions by

$$K_\nu(x) = \left(\frac{\pi}{2x}\right)^{1/2} W_{0,\nu}(2x).$$

Corollary 3.4. Suppose $x \in \mathbb{C}$, $|\arg x| < \pi$, $t > 0$, $\nu \in \mathbb{R}$. Then
(3.3)

$$e^{2t\Re x} |K_\nu(tx)|^2 = \pi(2|x|)^{2\nu} \int_0^\infty e^{-tu} |2x+u|^{-2\nu-1} {}_2F_1\left(\nu + \frac{1}{2}, \nu + \frac{1}{2}; 1; \frac{u(4\Re x + u)}{|2x+u|^2}\right) du.$$

Corollary 3.5. Suppose $x \in \mathbb{C}$, $|\arg x| \leq \frac{1}{2}\pi$, $\nu \in \mathbb{R}$ and $\tau \geq 1$. Then

$$(3.4) \quad \left| \frac{K_\nu(\tau x)}{K_\nu(x)} \right|^2 \leq e^{(1-\tau)2\Re x}.$$

For $\nu \in \mathbb{R}$, let $z = p(\nu)$ denote the largest negative zero of the hypergeometric function

$${}_2F_1\left(\nu + \frac{1}{2}, \nu + \frac{1}{2}; 1; z\right)$$

if it exists; otherwise set $p(\nu) = -\infty$.

Corollary 3.6. Suppose that $\nu \in \mathbb{R}$, $x \in \mathbb{C}$ such that $|\arg x| < \pi$ if $p(\nu) = -\infty$, and otherwise $|\arg x| \leq \theta$, where θ is the angle between $\frac{1}{2}\pi$ and π for which

$$\tan \theta = -\frac{1}{(-p(\nu))^{1/2}}.$$

Then the function

$$(3.5) \quad f(t) = e^{2t\Re x} |K_\nu(tx)|^2$$

is (strictly) completely monotone on $(0, \infty)$.

Corollary 3.7. Suppose that $-\frac{1}{2} \leq \nu \leq \frac{1}{2}$, and $x \in \mathbb{C}$, $|\arg x| < \pi$. Then the function (3.5) is (strictly) completely monotone on $(0, \infty)$.

Erdélyi [2, page 875] mentioned that the hypergeometric function appearing in connection with Bessel functions can be expressed as a Legendre function by means of

$${}_2F_1(a, a; 1; 1-z) = z^{-\frac{1}{2}a} P_{-a}\left(\frac{2}{z} - 1\right).$$

If $\nu \in \mathbb{R}$ and $|\arg x| \leq \frac{1}{2}\pi$ then the function

$$t \mapsto |K_\nu(tx)|^2$$

is also completely monotone on $(0, \infty)$ because the product of completely monotone functions is again completely monotone.

Example: Consider $\nu = 2$. By numerical calculation we find $p(2) \approx -0.4573617040$. The function

$$t \mapsto e^{2t\Re x} |K_2(tx)|^2, \quad t > 0$$

is completely monotone provided that $|\arg x| \leq \theta$, where

$$\theta \approx 2.165428404.$$

Boris Belinskiy also asked questions involving the Whittaker function $M_{k,m}$. For example, one would like to estimate the quotient

$$\frac{M_{k,m}(z)}{M_{k,m}(\tau z)}$$

when k is purely imaginary, m is real and $|\arg z| < \frac{1}{2}\pi$. However, there does not seem to exist a formula for $M_{k,m}$ analogous to (2.1). Therefore, different methods would have to be used for the Whittaker function $M_{k,m}$.

REFERENCES

- [1] H. Buchholz. *Die konfluente hypergeometrische Funktion mit besonderer Berücksichtigung ihrer Anwendungen*. Ergebnisse der angewandten Mathematik. Bd. 2. Springer-Verlag, Berlin, 1953.
- [2] A. Erdélyi. Integral representations for products of Whittaker functions. *The London, Edinburgh and Dublin philosophical magazine and journal of science* 26, 871–877 (1938).
- [3] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, editors. *NIST handbook of mathematical functions*. Cambridge University Press, Cambridge, 2010.
- [4] F. W. J. Olver. *Asymptotics and Special Functions*. Academic Press, New York 1974.
- [5] D. Widder. *The Laplace Transform*. Princeton University Press, Princeton, New Jersey, 1941.

DEPARTMENT OF MATHEMATICAL SCIENCES,, UNIVERSITY OF WISCONSIN–MILWAUKEE,
P. O. BOX 413,, MILWAUKEE, WI 53201, U.S.A.